

» Print

This copy is for your personal, non-commercial use only. To order presentation-ready copies for distribution to colleagues, clients or customers, use the Reprints tool at the top of any article or visit: www.reutersreprints.com.

Scientists warn geoengineering may disrupt rainfall

Wed, Jun 6 2012

By Chris Wickham

LONDON (Reuters) - Large-scale engineering projects aimed at fighting global warming could radically reduce rainfall in Europe and North America, a team of scientists from four European countries have warned.

Geoengineering projects are controversial, even though they are largely theoretical at this point. They range from mimicking the effects of large volcanic eruptions by releasing sulphur dioxide into the atmosphere, to deploying giant mirrors in space to deflect the sun's rays.

Proponents say they could be a rapid response to rising global temperatures but environmentalists argue they are a distraction from the need to reduce man-made carbon emissions.

Critics also point to a lack of solid research into unintended consequences and the absence of any international governance structure for such projects, whose effects could transcend national borders.

A small geoengineering experiment in the UK was recently abandoned due to a dispute over attempts by some of the team involved to patent the technology.

In this new study scientists from Germany, Norway, France and the UK used four different computer models that mimic the earth's climate to see how they responded to increased levels of carbon dioxide coupled with reduced radiation from the sun.

Their scenario assumed a world with four times the carbon dioxide concentration of the preindustrial world, which lead author Hauke Schmidt says is at the upper end, but in the range of what is considered possible at the end of this century.

They found that global rainfall was reduced by about 5 percent on average using all four models.

"Climate engineering cannot be seen as a substitute for a policy pathway of mitigating climate change through the reduction of greenhouse gas emissions," they said in the study, published in *Earth System Dynamics*, an open access journal of the European Geosciences Union.

Under the scenario studied, rainfall diminished by about 15 percent, or about 100 millimeters per year, compared to pre-industrial levels, in large areas of North America and northern Eurasia.

Over central South America, all the models showed a decrease in rainfall that reached more than 20 percent in parts of the Amazon region.

(Editing by Mark Heinrich)

© Thomson Reuters 2011. All rights reserved. Users may download and print extracts of content from this website for their own personal and non-commercial use only. Republication or redistribution of Thomson Reuters content, including by framing or similar means, is expressly prohibited without the prior written consent of Thomson Reuters. Thomson Reuters and its logo are registered trademarks or trademarks of the Thomson Reuters group of companies around the world.

Thomson Reuters journalists are subject to an Editorial Handbook which requires fair presentation and disclosure of relevant interests.

This copy is for your personal, non-commercial use only. To order presentation-ready copies for distribution to colleagues, clients or customers, use the Reprints tool at the top of any article or visit: www.reutersreprints.com.