

« back to **CNN**health.com

Patient Care

**Health Information**



For Medical Professionals

Research

Education

Request an Appointment

Find a Doctor

Find a Job

Log in to Patient Account

Enter search term here

SEARCH

Diseases and Conditions

Symptoms

Drugs and Supplements

Tests and Procedures

Healthy Lifestyle

First Aid

Home Diseases and Conditions Radiation sickness Basics Symptoms

# Radiation sickness

Text Size:

Print Share Reprints

Basics

Resources

Definition

Symptoms

Causes

Tests and diagnosis

Complications

Treatments and drugs

Free

E-Newsletters

Subscribe to receive the latest updates on health topics.

About our newsletters

- Housecall
- Alzheimer's caregiving
- Living with cancer

Enter e-mail

SUBSCRIBE

RSS Feeds

Get **free personalized** health guidance for you and your family.

GET STARTED

## Symptoms

By Mayo Clinic staff

The severity of signs and symptoms of radiation sickness depends on how much radiation you've absorbed. How much you absorb depends on the strength of the radiated energy and the distance between you and the source of radiation.

### Absorbed dose and duration of exposure

The absorbed dose of radiation is measured in a unit called a gray (Gy). Diagnostic tests that use radiation, such as an X-ray, result in a small dose of radiation — typically well below 0.1 Gy, focused on a few organs or small amount of tissue.

Signs and symptoms of radiation sickness usually appear when the entire body receives an absorbed dose of at least 1 Gy. Doses greater than 6 Gy to the whole body are generally not treatable and usually lead to death within two days to two weeks, depending on the dose and duration of the exposure.

### Initial signs and symptoms

The initial signs and symptoms of treatable radiation sickness are usually nausea and vomiting. The amount of time between exposure and when these symptoms develop is an indicator of how much radiation a person has absorbed.

After the first round of signs and symptoms, a person with radiation sickness may have a brief period with no apparent illness, followed by the onset of new, more serious symptoms.

In general, the greater your radiation exposure, the more rapid and more severe your symptoms will be.

#### Early symptoms of radiation sickness

	Mild exposure (1-2 Gy)	Moderate exposure (2-6 Gy)	Severe exposure (6-8 Gy)	Very severe exposure (8-10 Gy or higher)
Nausea and vomiting	Within 6 hours	Within 2 hours	Within 1 hour	Within 10 minutes
Diarrhea	--	Within 8 hours	Within 3 hours	Within 1 hour
Headache	--	Within 24 hours	Within 4 hours	Within 2 hours
Fever	--	Within 3 hours	Within 1 hour	Within 1 hour

#### Later symptoms of radiation sickness

Dizziness and disorientation	--	--	Within 1 week	Immediate
Weakness, fatigue	Within 4 weeks	Within 1-4 weeks	Within 1 week	Immediate

Advertisement

**MAYO CLINIC WELLNESS SOLUTIONS FOR Fibromyalgia**

- Integrative medicine
- Gentle yoga
- Daily nutrition guide

TAKE CHARGE OF YOUR HEALTH **ACT NOW** >>>

Check out these best-sellers and special offers on books and newsletters from Mayo Clinic.

Try Mayo Clinic Health Letter free!

Mayo Clinic Wellness Solutions for Fibromyalgia — DVD

[Heartburn Symptoms](#)

Get Information On Heartburn

The Mayo Clinic Diet — stay connected with us on Facebook!

[Cell Phone Radiation](#)

Shields 99.95% of Radiation Get

RadiGuard Protection Today

[RadiGuard.com](#)

[Rheumatoid Pain Symptoms](#)

Leading the way in joint pain info! Get rheumatoid pain symptoms info.

[ArthritisResolutions.com](#)

Ads by Google

Advertising and sponsorship policy  
Advertising and sponsorship opportunities

healing, low  
blood  
pressure

[About this site](#)

[Site help](#)

[Contact us](#)

[E-newsletter](#)

[Blogs](#)

[Podcasts](#)

[Slide shows](#)

[Videos](#)

[Site map](#)

[Reprints and permissions](#)

Source: Adapted from "Bushberg JT. Radiation exposure and contamination. The Merck Manuals: The Merck Manual for Healthcare Professionals" and "Upton AC. Radiation injury. In: Goldman L, et al., eds. Cecil Medicine. 23rd ed. Philadelphia, Pa.: Saunders Elsevier; 2007."

### When to see a doctor

An accident or attack that causes radiation sickness would no doubt cause a lot of attention and public concern. If an event occurs, monitor radio, television or online reports to learn about emergency instructions for your area.

If you know you've been exposed to radiation, seek emergency medical care.

## Definition

## Causes

## References

May 8, 2010

DS00432

© 1998-2011 Mayo Foundation for Medical Education and Research (MFMER). All rights reserved. A single copy of these materials may be reprinted for noncommercial personal use only. "Mayo," "Mayo Clinic," "MayoClinic.com," "EmbodyHealth," "Enhance your life," and the triple-shield Mayo Clinic logo are trademarks of Mayo Foundation for Medical Education and Research.

Find Mayo Clinic on

[Facebook](#)

[Twitter](#)

[YouTube](#)

[Privacy policy](#) (Updated Nov. 5, 2010)

[Terms and conditions of use policy](#) (Updated July 8, 2010)

LEGAL CONDITIONS AND TERMS OF USE APPLICABLE TO ALL USERS OF THIS SITE. ANY USE OF THIS SITE CONSTITUTES YOUR AGREEMENT TO THESE TERMS AND CONDITIONS OF USE.

© 1998-2011 Mayo Foundation for Medical Education and Research. All rights reserved.



We comply with the [HONcode standard](#) for trustworthy health information: [verify here](#).