

**Navigation**

[UCBNE Home Page](#)
[Overview](#)
[Undergraduate](#)
[Graduate](#)
[News and Events](#)
[People](#)
[Research](#)
[Services](#)
[Careers](#)
[Class Schedule](#)
[Fall 2011 Class Schedule](#)
[Spring 2011 Class Schedule](#)
[NE Courses and Websites](#)
[create content](#)
[Public Forums](#)
[Colloquiums](#)
[ANS Berkeley Student Section](#)

Home**UCB Food Chain Sampling Results**

[Back to Main UCB Air and Water Sampling Page](#)

Radionuclides, once deposited by rainwater or air onto the ground, will find their way through the ecosystem. We are already tracking its path from [rainwater](#) to [creek runoff](#) to [tap water](#), but we would also like to monitor how much these isotopes that make their way into our food. For example, how much gets taken up by the grass and eventually winds up in our [milk](#)?

We have been collecting produce that is as local as possible to test for the radioactive isotopes. We might expect different kinds of plants to take up different quantities of cesium and iodine, so we are trying to measure as many different plants and fruits as we are able to. So far, we have measured [spinach](#), [strawberries](#), [cilantro](#), [grass](#), and [mushrooms](#). We have also measured local [topsoil](#).

In the tables below, we are providing two numbers for each of the isotopes. The first is a standard concentration unit of Becquerel per kilogram (Bq/kg) which is the number of particles decaying per second in each kilogram of the sample. The number in parentheses after the activity is the number of kilograms that one would need to consume to equal the radiation exposure of a single round trip flight from San Francisco to Washington D.C. (0.05 mSv). For more information on how this equivalent dose is calculated, the details are here: [How Effective Dose is Calculated](#)

The experimental setup used for the food testing is the same setup used for the [Rainwater Collection Experiment](#).

Topsoil

source: Alameda, CA

Collection Date	Sample Mass	I131	I132	Cs134	Cs137	Te132	Data
	kg	Bq/kg (kg**)	Bq/kg (kg**)	Bq/kg (kg**)	Bq/kg (kg**)	Bq/kg (kg**)	
4/6/2011	1.04	12.5 ± 0.4 (89)	Less than MDA	0.99 ± 0.13 (2,610)	1.5 ± 0.2 (2,510)	Less than MDA	data

Grass

source: Alameda, CA

Collection Date	Sample Mass	I131	I132	Cs134	Cs137	Te132	Data
	kg	Bq/kg (kg**)	Bq/kg (kg**)	Bq/kg (kg**)	Bq/kg (kg**)	Bq/kg (kg**)	
4/3/2011 10:00	0.400	9.8 ± 0.5 (113)	Less than MDA	6.9 ± 0.3 (373)	6.9 ± 0.3 (537)	Less than MDA	data
4/5/2011 08:30	0.360	6.9 ± 0.8 (161)	Less than MDA	4.8 ± 0.4 (542)	3.1 ± 0.6 (1,195)	Less than MDA	data

Wild Mushrooms

source: Alameda, CA

Collection Date	Sample Mass	I131	I132	Cs134	Cs137	Te132	Data
	kg	Bq/kg (kg**)	Bq/kg (kg**)	Bq/kg (kg**)	Bq/kg (kg**)	Bq/kg (kg**)	
4/2/2011 10:00	0.390	8.4 ± 0.5 (132)	Less than MDA	0.63 ± 0.19 (4,110)	0.47 ± 0.19 (7,830)	Less than MDA	data

Spinach

"Best By" Date	Food Mass	I131	I132	Cs134	Cs137	Te132	Data
	kg	Bq/kg (kg**)	Bq/kg (kg**)	Bq/kg (kg**)	Bq/kg (kg**)	Bq/kg (kg**)	
4/8/2011	0.284	2.8 ± 0.8 (403)	Less than MDA	Less than MDA	Less than MDA	Less than MDA	data

Strawberries

Collection Date	Food Mass	I131	I132	Cs134	Cs137	Te132	Data
	kg	Bq/kg (kg ^{**})	Bq/kg (kg ^{**})	Bq/kg (kg ^{**})	Bq/kg (kg ^{**})	Bq/kg (kg ^{**})	
4/1/2011	1.23	0.25 ± 0.14 (4,370)	Less than MDA	0.69 ± 0.07 (3,780)	0.67 ± 0.10 (5,520)	Less than MDA	data

Cilantro

Collection Date	Food Mass	I131	I132	Cs134	Cs137	Te132	Data
	kg	Bq/kg (kg ^{**})	Bq/kg (kg ^{**})	Bq/kg (kg ^{**})	Bq/kg (kg ^{**})	Bq/kg (kg ^{**})	
4/4/2011	0.500	Less than MDA	Less than MDA	Less than MDA	Less than MDA	Less than MDA	data

** The number in parentheses is the number of kilograms of the item that one would need to consume to equal the radiation exposure of a single round trip flight from San Francisco to Washington D.C. (0.05 mSv). To see how we calculate these numbers, please visit [our explanation of the equivalent dose calculation](#).

Note: "MDA" is the estimated minimum detectable activity for a given isotope in the detector.