

Saturday
12th June 2004
[Text only](#)

New visitors: [Create your BBCi membership](#)

[BBC Homepage](#)

[Add Article](#)
[Search 360](#)

[Feedback](#)
[360 FAQ](#)
[About 360](#)
[Site News](#)
[Help!](#)

[Latest](#)
[Discussions](#)

[360](#)
[Community](#)

[The Book of](#)
[Solutions](#)

[360 Missions](#)

Site index:

[*ABCDEF](#)
[GHIJKLM](#)
[NOPQRS](#)
[TUVWXYZ](#)

Search 360:

[About the BBC](#)

[Contact Us](#)

[Help](#)

Like this page?
[Send it to a friend!](#)

RESOLVING WATER CRISIS BY INTERLINKING RIVERS AND CLOUD SEEDING



The basic foundation of Indian economy is agriculture on which more than two thirds of the population depend for their livelihood and two-thirds of the labour force are employed in it. Although there was a drought in 1987, there was food production of 135 million tonnes (M.T.) of which 50 M.T. was due to irrigation under the major reservoirs completed in the post independent era. Although the country was importing food grains till the mid 1960s food security was achieved later due to introduction of high yielding seed varieties for wheat, rice and other cereals and application of fertilisers and crop protection technologies.

Food grain production increased four-fold from 50 M.T. in 1950-51 to about 200 M.T. today, leading to augmentation of per capita availability of food grains from 400 to about 500 grams and during the period while the land under agriculture rose from 50 million hectares (M.ha) to about 128 million hectares. With the same acreage under wheat as in USA, India emerged as the second largest after China with 72 M.T. In 1995 World Watch Institute warned that population rich but land-hungry countries like China and India will have to resort to substantial food import in another 30 years mainly due to fast decreasing per capita availability of arable land and irrigation water. China, USA and India are producing nearly 400, 320, and 200 million tons of food grains by utilising about 90, 190 and 188 million hectares of land. It clearly shows that Indian agriculture has great scope for development and for this purpose irrigation facilities must be created on large scale and new varieties of high yielding crops must be developed by utilising the advances in Biotechnology.

Our population has already touched 1000 billions and it is going to touch 1350 billions by 2025 AD and 1650 billions by 2050 AD. In order to feed this exploding population we have to increase the food production to 325 million tonnes by 2025 AD and 450 million tonnes by 2050 AD. In particular the production of rice must be increased from the present annual production of 85 million tonnes to 135 million tonnes by 2025 AD and 200 million tonnes by 2050 AD.
<http://www.geocities.com/jvmnaidu/cloudseeding.html>

The water use for different purposes upto 2050 AD are estimated as follows.(Table –1)

The present utilisable surface and ground water is estimated at 1122 billion cu.m. and consequently there is bound to be a water deficit of 126 billion cu.m. by 2025 AD and 534 billion cu.m. by 2050 AD River water storages must be increased to overcome the difficulties caused by frequent failures of monsoons and also to meet the ever growing demands of summer seasons. The present live storages of about 174 billion cu.m. constitute about 10%. Of the total water potential available in the country. Even if a low estimate of future water requirements of 1300 billion cu.m. is to be met, a storage of 600 billion cu.m. is required. (See Table 2 and 3). In order to maintain minimum summer flows to dilute the impact of treated waste

water discharges from municipalities and industries, additional storage is needed. Further 50 billion cu.m. is needed to maintain ecological balances in water courses. In addition to the above mentioned live storage the projects under construction and those under consideration envisage additional storages of 75 billion cu.m. and 130 billion cu.m. respectively. Thus there is an urgent need to conserve river waters by transferring the surplus flows from rivers like Ganges, Mahanadi and Godavari to water deficit rivers like Krishna, Pennar and Cauvery. Environmentalists predict that unless more land is brought under agriculture and more reservoirs are constructed across the rivers to store water that is wasted into the sea there will be water famine and food famine in India in general and the Southern states in particular. (See Table 4)

<http://www.geocities.com/jvmnaidu/cloudseeding.html>

GANGA-CAUVERY WATER LINK:

<http://www.indianest.com/voices/v002.htm>

<http://www.tribuneindia.com/2003/20030313/science.htm#1>

<http://www.ficci.com/ficci/march5-rivers-goswami.htm>

<http://rrtd.nic.in/LINKING%20RIVERS.htm>

According to the Indian Water Resources Society “A note on the National Water Grid prepared by the then Central Water Commission with 3 possible alignments for Ganga-Cauvery links along with other links were brought out. Further studies were made by Dr.K.L.Rao who advocated one of the alignments for Ganga-Cauvery Link along with a few other links including the Brahmaputra and Ganga Link. The 2640 km long Ganga-Cauvery link essentially envisaged the withdrawal of 1680 cumecs (60,000 cusecs) of the flood flows of the Ganga near Patna for about 150 days in a year and pumping about 1400 cumecs (50,000 cusecs) of this water over a head of 5409 meters (1800ft) for transfer to the Peninsular region and utilising the remaining 280 cumecs (10,000 cusecs) in the Ganga basin itself. The proposal envisaged utilisation of 2.59 million hectare meters of Ganga waters to bring under irrigation and additional area of 4 M.Ha. Dr.Rao had also proposed a few additional links like (a) Brahmaputra-Ganga link to transfer 1800 to 3000 cumecs with a lift of 12 to 15m (b) Link transferring 300 cumecs of Mahanadi waters southwards (c) Canal from Narmada to Gujarat and Western Rajasthan with a lift of 275 m and (d) links from rivers of the Western ghats towards East. Dr.Rao had estimated his proposals to cost about Rs.12,500 crores. Very roughly at 1995 prices the Ganga Cauvery link alone would amount to about Rs.70,000 crores (capital cost). The annual costs including cost of power would be around Rs.30,000 per ha. As further seen, the present NWDA proposals for inter linking river between Ganga and Cauvery at present prices would cost on around Rs.15,000 per ha. Annually. The proposals were examined by the Central Water Commission and found to be grossly under-estimated. It was also observed that the scheme would require large blocks of power (5 to 7 million kw) for lifting water. It will also have no flood control benefits. Therefore, the proposal was not pursued as such.

N.W.D.A.PROPOSAL: National Water Development Agency Proposes

1. To provide terminal storages on Mahanadi and Godavari to transfer a surplus water by gravity and lift to the drought-prone areas of Andhra Pradesh, Tamil Nadu and other states.
2. To plan for diversion of water from the West flowing rivers of Kerala and the East flowing rivers of Tamil Nadu.
3. To plan for construction of small storage dams and to inter link the rivers flowing on the west coast for transferring water to the East of the Western ghats through small tunnels or mountain passes and thereby augment the water yield in the rivers of the Southern states.

Surplus in Mahanadi is 22000 Mm³ (770 TMC) and Godavari is 28000 Mm³ (1000 TMC) against the present assessment of 11500Mm³(400 TMC) and 15000 Mm³ (530 TMC) respectively.

I. From Manibhadra dam on Mahanadi 8000 Mm³ equivalent to 280 TMC is proposed to be transferred through Mahanadi-Godavari link to deliver 6500 Mm³ (230TMC) into Godavari at Rajahmundry.

II. GODAVARI TO KRISHNA: Surplus in Godavari is calculated at 21500 Mm³ (15000+6500) (760 TMC) and this is proposed for transfer into Krishna through 3 links.

(a) 1200 Mm³ (42 TMC) will be diverted through Polavaram-Vijayawada link to supplement needs of the delta.

(b) 4370 Mm³ (154 TMC) is proposed to be transferred through Inchampalli-Pulichintala link totake over part of the command under Nagarjuna Sagar left and right bank canals as exchange.

(c) 14000 Mm³ (500 TMC) of water is proposed to be discharge into Nagarjuna Sagar reservoir after considering needs of enroute irrigation and transmission losses. The electrical power requirements for lifting the water to Pulichintala and Nagarjuna Sagar will be about 110 MW and 1650 MW respectively.

III. KRISHNA TO PENNAR: Out of 14000 Mm³ discharged into Nagarjuna Sagar 12000 Mm³ (420 TMC) is proposed for diversion through Nagarjuna Sagar-Somasila link and the balance is utilized under part command of sagar left canal ir exchange. After considering enroute irrigation under Sagar right canal in exchange and irrigation under other canals about 9800 Mm³ (346 TMC) reaches Somasila. Since this link provides water on exchange basis to Sagar canals an equivalent quantity of water may be diverted from Srisailam and Alamatti to other needy areas. 2300 Mm³ (80 TMC) is proposed to be diverted from Srisailam through Srisailam-Proddutur link to reach the barrage at Proddutur. About 2000 Mm³ (70 TMC) is proposed for diversion from Alamatti through Alamatti-Pennar link to cate for enroute irrigation under Krishna and Pennar basins.

IV. PENNAR TO CAUVERY: About 9500 Mm³ (335 TMC) of water is proposed to be transferred through Pennar-Cauvery link into Grand Anicut deducting for enroute irrigation and Madras water supply about 5000Mm³ (176 TMC) will reach Grand Anicut and out of this water about 3000 Mm³ (106 TMC) will be used in Cauvery delta.

V. CAUVERY TO VAIGAI: Out of 5000Mm³ reaching Upper Anicut, about 2000 Mm³ (70 TMC) is proposed to be diverted through Cauvery-Vaigai link for utilization in Cauvery, Vaigai and other stream in between Vaigai and Vaippar.

ECONOMICS : Peninsular link project from Mahanadi to Cauvery costs Rs.50,000 crores for main works and additional amount of Rs.30,000 crores will be needed for branchcanals, field channels.

AUGMENTING ANNUAL YIELD IN RIVERS AND LAKES BY CLOUD SEEDING

The water droplets in the clouds are so small (10 micro-meters in radius) that the cannot reach the earth without evaporating. Hence if large rain drops (500 micro-meters in radius) tht fall on the earth without evaporating are to be formed, about a million small cloud droplets must coagulate since mere adhesion and condensation are not enough. There are 2 theories about the mechanics of rain-drop formation, each of which is valid under different climates.

The first theory of rain-drop formation in warm clouds (do not reach freezing levels) is based on "Langmuir chain-reaction" that mostly occurs in the hot, humid atmosphere of the tropics. In a rising cloud, the up-currents hold the big-drops in suspension while the small ones are caught up and merge with the big ones. If the large drops become too big, they burst into fragments that again collide and coalesce with small droplets and ultimately fall to earth as rain-drops. The second

theory for cold clouds (tops touch freezing levels) is based on "Bergeron chain-reaction". The clouds that precipitate consist at the higher levels, of ice crystals. For the formation of big droplets in the cloud an ice-phase is a precondition. For transition from the liquid to solid phase a freezing nucleus is essential. The freezing nuclei allow supercooling of water droplets upto 15o C before ice-formation occurs. Infact pure water suspended in air does not freeze until it touches about minus 40o C temperature. The ice-crystals grow rapidly by absorbing all the supercooled droplets on their way down and unwieldy lumps of ice are formed and they melt a about zero degree centigrade to become rain-drops that fall on earth.

In w arm clouds to be seeded there is a serious shortage of large wt er droplets over 20 microns in diameter and this reduces the efficiency of the cloud to give rain. In order to induce such clouds to give rain the coalescence mechanisms have to be stimulated by injecting the clouds with the hydrosopic sodium chloride particles from a small aircraft. The Langmuir chain-reaction gets a boost and the clouds which were formerly unproductive become productive. Similarly the dearth of ice-crystals in cold clouds is removed by seeding them with silver iodide to ensure one nucleus per litre of cloud air so that such seeding generates additional ice-crystals and thereby accelerates the rain-forming processess.

CLOUD SEEDING FOR ARTIFICIAL RAINS: Since salt crystals are abundant in oceanic regions they favour larger cloud droplets that collide and coalescence to initiate rain fall with in the life time of the clouds. But the atmosphere over continental regions contain much smaller and more numerous condensation nuclei and hence the medium sized clouds formed in such regions usually dissipate before coalescence mechanism has had a chance to initiate rain. When right conditions exist, the cumulus clouds can be stimulated to grow larger and last longer. The freezing of millions of droplets in the cloud release enomous latent heat energy and that makes the cloud buoyant, causing it to grow larger, thereby efficiently processing more wter for a longer time period than would have occurred without the cloud-seeding process.

COLD-CLOUD SEEDING: Introduction of massive dosages of silver iodide into the clouds at minus 10o C at the rate of 100 to 1000 nuclei per liter of cloud air is known as dynamic seeding. In this process the conversion of super-cooled water i the cloud into ice-crystals (though Bergeron-Findesen reaction mechanism) releases latent heat that increases buoyancy, thereby increasing the size of the cloud with better organisation of low level inlows with the probability of cloud-merger and area wise enhancement of rain-fall. Dynamic seeding of individual clouds on a random basis in USA, indicated significant increase in rain-fall by a factor of three.

REFERENCES – WEB SITES ON CLOUD SEEDING

<http://www.bbc.co.uk/dna/360/A811559>

<http://www.bbc.co.uk/dna/360/A787656>

http://www.rap.ucar.edu/general/press/presentations/wxmod_overview/index.htm

<http://twri.tamu.edu/twripubs/WtrResrc/v20n2/>

<http://www.xmission.com/~nawc/wm.html>

<http://www.license.state.tx.us/weather/summary.htm>

<http://www.edwardsaquifer.net/cloudseeding.html>

<http://physicsweb.org/article/world/11/5/3>

<http://www.countyofsb.org/pwd/water/cloud.htm>

<http://www.physics.isu.edu/weather/kmddbdc/cloudseed.pdf>

<http://www.geocities.com/CapeCanaveral/Lab/5833/seed/seed1.html>

<http://www.aoml.noaa.gov/hrd/nhurr97/CSEED.HTM>

<http://www.ghcc.msfc.nasa.gov/GOES/globalir.html>

<http://www.dar.csiro.au/publications/cloud.htm>

http://dir.yahoo.com/Science/Earth_Sciences/Meteorology/Weather_Modification/

TABLE-1: WATER USE FOR DIFFERENT PURPOSES (in billion cu.m.)

S.No.	Purpose	2000	AD 2025	AD 2050	AD
1.	Drinking Water	44	77	93	
2.	Irrigated Agriculture	520	909	1072	
3.	Hydro power	27	70	212	
4.	Industries	30	120	200	
5.	Ecology, Recreation etc.	41	72	80	
	TOTAL	662	1248	1656	

TABLE-2: PRODUCTION OF FOOD GRAINS IN INDIA

Crop	1950-51	60-61	70-71	80-81	90-91	97-98	2010	2025	2050	Annual growth rate		
	94-2010	2010-2025	2025-2050									
Rice	20	34	42	54	74	82	104	134	198	2.2	1.7	1.6
Wheat	7	11	24	37	55	66	80	110	170	2.6	2.1	1.8
Coarse cereals	15	24	31	29	33	34						
Pulses	8	13	12	11	15	13	15	21	40	3.0	2.6	2.6
Total food grains	50	80	110	130	177	193	220	287	435	2.2	1.8	1.8

All the figures are in Million tons.

TABLE: 3 FOOD-SHORTAGE FORECAST IN INDIA FOR 2025 A.D. AND 2050 A.D.

S.No.	Particulars	1950	1975	2000	2025	2050
1.	Population, m.	360	640	1000	1350	1650
2.	Land Area, M.ha.	22	55	90	150	200
3.	Food grains, M.t.	50	120	195	325	450
4.	Rice, M.t.	22	50	85	135	200
5.	Wheat, M.t.	7	28	66	110	170
6.	Irrigation water B.cum	100	250	400	640	900

TABLE:4 FOOD SHORTAGE IN SOUTH INDIAN STATES (1991-2050)
Population, Rice Demand, Production and Deficit

State	1991	2000	2025	2050
	Pop	dem	Prod	def
Andhra Pradesh	66	13	11	2
Tamil Nadu	55	11	5.5	5.5
Karna-taka	49	10	2.5	7.5
Kerala	30	6	1	5
Total	200	40	20	20

NOTE: Population (in Millions)

Rice production, Demand and shortages (in Million tons).

Pop(Population), dem(Demand), Prod(Production), def(Deficit)

Start a New Discussion

If you [register](#) you can discuss this Article with other Members.

[Terms of Use](#) | [Privacy & Cookies Policy](#)